Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently Amended) A method of coating a cell producing a very thin uniform crosslinked hydrocolloid coating of a single cell produced by capillary means comprising the steps of:

placing the cell in a solution of hydrocolloid;

removing the cell from the solution of hydrocolloid by sucking the cell into a capillary; and

placing the cell in a cross-linking solution after removing the cell from the solution of hydrocolloid, thereby providing the cell with a thin coating with a thickness of 1 to 5% of the cell diameter, in cases of iota-carrageenan and kappa-carrageenan and between 6 to 8% of the of the cell diameter in cases of LMP and alginate of the hydrocolloid; and storing the cell in solution.

- 2. (Original) A method as defined in Claim 1, wherein the hydrocolloid is an alginate.
- 3. (Previously Presented) A method as defined in Claim 1, wherein the hydrocolloid is Na-alginate.

- 4. (Previously Presented) A method as defined in Claim 1, wherein the hyrocolloid is low-methoxy pectin (LMP).
- 5. (Previously Presented) A method as defined in Claim 1, wherein the hydrocolloid is either κ or ι carrageenan.
- 6. (Previously Presented) A method as defined Claim 1, wherein the hydrocolloid solution is in Calcium Adjusted Modified Marc's Ringer (CAMMR) solution.
- 7. (Previously Presented) A method as defined in Claim 1, wherein the cell is a *Xenopus laevis* egg and embryos.
- 8. (Previously Presented) A method as defined in Claim 1, wherein the cross-linking solution is a solution of Ca, Ba or K ions.
- 9. (Original) A method as defined in Claim 8, wherein the cross-linking solution is a solution of CaCl₂, BaCl₂ or KCl.
- 10. (Previously Presented) A method as defined in Claim 9, wherein the cross-linking solution of CaCl₂ or BaCl₂ is at a concentration of from 0.25% and the KCl solution is at a concentration of 0.5%.

Claims 11-12 (Cancelled)

- 13. (Previously Presented) A method as defined in Claim 1, wherein the alginate has a high mannuronic acid (M) content.
- 14. (Previously Presented) A method as defined in Claim 13 wherein the mannuronic acid (M) content of the alginate is from about 29 to about 61 %.

Claims 15-20 (Cancelled)

- 21. (New) A method as defined in claim 1, where the cell or embryo to be coated is maneuvered by a thin capillary having an approximate or smaller diameter than the diameter of the cell, in such a manner that the coating is forced to perform with a minimal thickness and volume.
- 22. (New) A method as defined in claim 1, where the coating is uniform on all sides of the coated cell.
 - 23. (New) A method as defined in claim 1, where the coating is transparent.
- 24. (New) A method as defined in claim 1, wherein the coating accumulates photons.
- 25. (New) A method as defined in claim 1, where the coating forms a microbial shield.

- 26. (New) A method as defined in claim 1, where the coating is resistant to hazardous material.
- 27. (New) A method as defined in claim 1, where the coating acts as an inhibitor against damage during freezing and thawing.
- 28. (New) A method of producing a very thin uniform cross-linked hydrocolloid coating of an embryo produced by capillary means comprising the steps of:

placing the embryo in a solution of hydrocolloid;

removing the embryo from the solution of hydrocolloid by sucking the embryo cell into a capillary; and

placing the embryo in a cross-linking solution after removing the embryo from the solution of hydrocolloid, thereby providing the embryo with a coating with a thickness of 1 to 5% of the embryo diameter, in cases of iota-carrageenan and kappa-carrageenan and between 6 to 8% of the embryo diameter in cases of LMP and alginate of the hydrocolloid; and

storing the embryo in solution.

29. (New) A method as defined in Claim 28, wherein the hydrocolloid is an alginate.

- 30. (New) A method as defined in Claim 28, wherein the hydrocolloid is Naalginate.
- 31. (New) A method as defined in Claim 28, wherein the hyrocolloid is low-methoxy pectin (LMP).
- 32. (New) A method as defined in Claim 28, wherein the hydrocolloid is either κ or ι carrageenan.
- 33. (New) A method as defined Claim 28, wherein the hydrocolloid solution is in Calcium Adjusted Modified Marc's Ringer (CAMMR) solution.
- 34. (New) A method as defined in Claim 28, wherein the cell is a *Xenopus laevis* egg and embryos.
- 35. (New) A method as defined in Claim 28, wherein the cross-linking solution is a solution of Ca, Ba or K ions.
- 36. (New) A method as defined in Claim 35, wherein the cross-linking solution is a solution of CaCl₂, BaCl₂ or KCl.
- 37. (New) A method as defined in Claim 36, wherein the cross-linking solution of CaCl₂ or BaCl₂ is at a concentration of from 0.25% and the KCl solution is at a concentration of 0.5%.

- 38. (New) A method as defined in Claim 28, wherein the alginate has a high mannuronic acid (M) content.
- 39. (New) A method as defined in Claim 38 wherein the mannuronic acid (M) content of the alginate is from about 29 to about 61 %.
- 40. (New) A method as defined in claim 28, where the cell or embryo to be coated is maneuvered by a thin capillary having an approximate or smaller diameter than the diameter of the cell, in such a manner that the coating is forced to perform with a minimal thickness and volume.
- 41. (New) A method as defined in claim 28, where the coating is uniform on all sides of the coated cell.
 - 42. (New) A method as defined in claim 28, where the coating is transparent.
- 43. (New) A method as defined in claim 28, wherein the coating accumulates photons.
- 44. (New) A method as defined in claim 28, where the coating forms a microbial shield.

- 45. (New) A method as defined in claim 28, where the coating is resistant to hazardous material.
- 46. (New) A method as defined in claim 28, where the coating acts as an inhibitor against damage during freezing and thawing.